Neuroscience Letters, 81 (1987) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abell, C.W. see Awasthi, Y.C., 159
Aceves, J., see Galarraga, E., 351
Agnati, L.F., see Fuxe, K., 183
Ajima, A. and Kato, T., Brain dialysis: detection of acetylcholine in the striatum of unrestrained and unanesthetized rats, 129
Akaike, N., see Tokutomi, N., 314
Amit, Z., see Ng Cheong Ton, J.M., 335
Amit, Z., see Schenk, S., 227
Angioy, A.M., Tomassini Barbarossa, I., Crnjar, R., Liscia, A. and Pietra, P., Reflex cardiax response to various olfactory stimuli in the blowfly, *Protophormia terraenovae*, 263
Aniksztejn, L., Roisin, M.P., Gozlan, H. and Ben-Ari, Y., Long-lasting potentiation produced by a phorbol ester in the hippocampus of the anaesthetized rat is not associated with a persistent enhanced release of excitatory amino acids, 291
Antal, A., see Barthó, L., 146
Armstrong, J.D., see Reiter, R.J., 171
Awasthi, Y.C., Singh, S.V., Shen, R.-s., Abell, C.W., Gessner, W. and Brossi, A., MPTP metabolites inhibit rat brain glutathione *S*-transferases, 159

Baile, C.A., see Marson, L., 35
Bandler, R., see Carrive, P., 273
Barden, N., see Gerstberger, R., 193
Bargas, J., see Galarraga, E., 351
Barthó, L., Pethö, G., Antal, A., Holzer, P. and Szolcsányi, J., Two types of relaxation due to capsaicin in the guinea pig isolated ileum, 146
Baudry, M., see Kessler, M., 221
Baudry, M., see Vodyanoy, V., 133
Bell, C., Mann, R. and Borri Voltattorni, C., Two classes of sympathetic nerves with different DOPA decarboxylase immunoreactivities exist in dog vas deferens, 19
Ben-Ari, see Aniksztejn, L., 291
Blähser, S., see Szabo, T., 245
Bloom, S.R., see Inyama, C.O., 111
Borri Voltattorni, C., see Bell, C., 19
Bowker, R.M., Evidence for the co-localization of somatostatin- and methionine-enkephalin-like immunoreactivities in raphe and gigantocellularis nuclei, 75
Bradford, H.F., Young, A.M.J. and Crowder, J.M., Continuous glutamate leakage from brain cells is balanced by compensatory high-affinity reuptake transport, 296
Bradford, H.F., see Docherty, M., 232
Britt, J.H., see Reiter, R.J., 171
Brossi, A., see Awasthi, Y.C., 159
Burt, D.R., see Sharif, N.A., 339

Carrive, P., Dampney, R.A.L. and Bandler, R., Exitation of neurones in a restricted portion of the midbrain periaqueductal grey elicits both behavioural and cardiovascular components of the defence reaction in the unanaesthetised decerebrate cat, 273
Caruso, A., see Prato, 303
Cassone, V.M., Roberts, M.H. and Moore, R.Y., Melatonin inhibits metabolic activity in the rat suprachiasmatic nuclei, 29
Cazalets, J.R., Nagy, F. and Moulins, M., Suppressive control of a rhythmic central pattern generator by an identified modulatory neuron in crustacea, 267
Clementi, G., see Prato, A., 303
Cortés, R., see Cymerman, U., 13
Crnjar, R., see Angioy, A.M., 263
Crowder, J.M., see Bradford, H.F., 296
Crutcher, K.A., see Saffran, B.N., 215
Cuénod, M., see Knöpfel, T., 188
Cymerman, U., Palacios, J.M., Cortés, R. and Skangiel-Kramska, J., Autoradiographic localization of muscarinic cholinergic receptors in visual areas of cat brain: variations in sensitivity of *N*-[^{3}H]methylscopolamine binding sites to carbachol and pirenzepine, 13

Damm, H., see Susilo, R., 325
Dampney, R.A.L., see Carrive, P., 273
Davis, C.J., see Inyama, C.O., 111
Deguchi, T., see Ichikawa, T., 24
Della-Fera, M.-A., see Marson, L., 35
Denizot, J.P., see Szabo, T., 245
Dietrichs, E., see Walberg, F., 82
DiPaolo, T., see Gerstberger, R., 193
Docherty, M., Bradford, H.F. and Wu, J.-Y., The preparation of highly purified GABAergic and cholinergic synaptosomes from mammalian brain, 232

Eneroth, P., see Forsberg, G., 151
Evans, R.J., see Fawthrop, D.J., 250
Evans, R.J., see Fawthrop, D.J., 257

Fagni, L., Zinebi, F. and Hugon, M., Helium pressure potentiates the *N*-methyl-D-aspartate- and D,L-homocysteate-induced decreases of field potentials in the rat hippocampal slice preparation, 285
Fawthrop, D.J. and Evans, R.J., Morphological changes in cultured astrocytes following exposure to calcium ionophores, 250
Fawthrop, D.J. and Evans, R.J., The morphological differentiation of cultured astrocytes induced by ionomycin: lack of dependence on protein kinase C activation, 257
Ferssiwi, A., see Villalobos, J., 89
Ferssiwi, A., see Villalobos, J., 95
Fischer-Colbrie, R., see Volknandt, W., 241
Forsberg, G., Wiesenfeld-Hallin, Z., Eneroth, P. and Södersten, P., Sexual behaviour induces nalaxone-reversible hypoalgesia in male rats, 151
Friedman, A. and Gutnick, M.J., Low-threshold calcum electrogenesis in neocortical neurons, 117
Fritzsch, B. and Wahnschaffe, U., Electron microscopical evidence for common inner ear and lateral line efferents in urodeles, 48
Fukushima, N., see Ueda, H., 309
Fuxe, K., Agnati, L.F., Merlo Pich, E., Meller, E. and Goldstein, M., Evidence for a fast receptor turnover of D_1 dopamine receptors in various forebrain regions of the rat, 183

Galarraga, E., Bargas, J., Martínez-Fong, D. and Aceves, J., Spontaneous synaptic potentials in dopamine-denervated neostriatal neurons, 351
Garzón, J., see Lérida, M., 155
Ge, M., see Ueda, H., 309
Gerstberger, R., DiPaolo, T. and Barden, N., Impaired regulation of neurohypophyseal vasopressin secretion in rats treated neonatally with monosodium-L-glutamate, 193
Gessner, W., see Awasthi, Y.C., 159
Gilbey, M.P., see Numao, Y., 279
Giuffrida, R., see Prato, A., 303
Goldstein, M., see Fuxe, K., 183
Gorman, K., see Schenk, S., 227
Gozlan, H., see Aniksztejn, L., 291
Gutnick, M.J., see Friedman, A., 117

Hirata, Y., see Ichikawa, T., 24
Höfle, G., see Susilo, R., 325
Hogan, C., see Westerman, R.A., 177
Holzer, P., see Barthó, L., 146
Hori, T., see Shioda, S., 53
Hughes, J., see Lo, W.W.Y., 331
Hugon, J., Vallat, J.M. and Leboutet, M.J., Cytotoxic properties of glutamate and aspartate in rat peripheral nerves: histological findings, 1
Hugon, M., see Fagni, L., 285
Hulsebosch, C.E., see McNeill, D.L., 57

Ichikawa, T., Ishida, I., Hirata, Y. and Deguchi, T., Monoclonal antibodies to monkey brain choline acetyltransferase: production and immunohistochemistry, 24
Ikeda, S.R., Schofield, G.G. and Weight, F.F., Somatostatin blocks a calcium current in acutely isolated adult rat superior cervical ganglion neurons, 123
Ikemoto, Y., see Tokutomi, N., 314
Inenaga, K., see Kasai, M., 319
Inyama, C.O., Wharton, J., Davis, C.J., Jackson, R.H., Bloom, S.R. and Polak, J.M., Distribution of vasoactive intestinal polypeptide binding sites in guinea pig genital tissues, 111
Ishida, , I., see Ichikawa, T., 24

Jackson, R.H., see Inyama, C.O., 111
Jacobson, K.A., see Jarvis, M.F., 69
Jarvis, M.F., Jacobson, K.A. and Williams, M., Autoradiographic localization of adenosine A_1 receptors in rat brain using [^{3}H]XCC, a functionalized congener of 1,3-dipropylxanthine, 69
Jones, R.S.G., Complex synaptic responses of entorhinal cortical cells in the rat to subicular stimulation in vitro: demonstration of an NMDA receptor-mediated component, 209

Kannan, H., see Kasai, M., 319
Kasai, M., Osaka, T., Inenaga, K., Kannan, H. and Yamashita, H., γ-Aminobutyric acid antagonist blocks baroreceptor-activated inhibition of neurosecretory cells in the hypothalamic paraventricular nucleus of rats, 319
Kato, T., see Ajima, A., 129
Kawai, N., see Pan-Hou, H., 199
Kessler, M., Baudry, M. and Lynch, G., Use of cystine to distinguish glutamate binding from glutamate sequestration 221
Knöpfel, T., Zeise, M.L., Cuénod, M. and Zieglgänsberger, W., L-Homocysteic acid but not L-glutamate is an endogenous *N*-methyl-D-aspartic acid receptor preferring agonist in rat neocortical neurons in vitro, 188

Koshiya, N., see Numao, Y., 279
Kramer, K., see Vodyanoy, V., 133
Kuba, K., see Tsuji, S., 139

Lacelle, G., see Schenk, S., 227
Land, P.W., see Simons, D.J., 100
Lauterio, T.J., see Marson, L., 35
Leboutet, M.J., see Hugon, J., 1
Lérida, M., Sánchez-Blázquez, P. and Garzón, J., Incidence of morphine withdrawal and quasi-abstinence syndrome in a model of chronic pain in the rat, 155
Liscia, A., see Angioy, A.M., 263
Lo, W.W.Y. and Hughes, J., Specific binding of [^{3}H]inositol trisphosphate to rat cerebral cortical microsomal membranes, 331
Lynch, G., see Kessler, M., 221
Lynch, G., see Vodyanoy, V., 133

Mann, R., see Bell, C., 19
Marson, L., Lauterio, T.J., Della-Fera, M.-A. and Baile, C.A., Immunohistochemical distribution of cholecystokinin, dynorphin A and Met-enkephalin neurons in sheep hypothalamus, 35
Martínez-Fong, D., see Galarraga, E., 351
McCulloch, J., see Reid, J., 165
McNeill, D.L. and Hulsebosch, C.E., Intraspinal sprouting of rat primary afferents after deafferentation, 57
Meller, E., see Fuxe, K., 183
Merlo Pich, E., see Fuxe, K., 183
Miller, J.A. and Zahniser, N.R., The use of ^{14}C-labeled tissue paste standards for the calibration of ^{125}I-labeled ligands in quantitative autoradiography, 345
Minota, S., see Tsuji, S., 139
Mizukawa, K., Ogawa, N., Sora, Y.H. and Sora, I., Alterations of the muscarinic cholinergic (mACh) receptors in the striatum of the MPTP-induced parkinsonian model in mice: in vitro quantitative autoradiographical analysis, 105
Mizuno, Y., Suzuki, K., Sone, N. and Saitoh, T., Inhibition of ATP synthesis by 1-methyl-4-phenylpyridinum ion (MPP^+) in isolated mitochondria from mouse brains, 204
Moore, R.Y., see Cassone, V.M., 29
Moulins, M., see Cazalets, J.R., 267
Muller, D., see Vodyanoy, V., 133
Mulligan, K.A. and Törk, I., Serotonergic axons form basket-like terminals in cerebral cortex, 7

Nagasawa, K., see Tokutomi, N., 314
Nagy, see Cazalets, J.R., 267
Nakai, Y., see Shioda, S., 53
Ng Cheong Ton, J.M. and Amit, Z., Effects of naloxazone on morphine–ethanol interaction in the conditioned taste aversion paradigm, 335
Nordby, T., see Walberg, F., 82
Numao, Y., Koshiya, N., Gilbey, M.P. and Spyer, K.M., Central respiratory drive-related activity in sympathetic nerves of the rat: the regional differences, 279

Ogawa, N., see Mizukawa, K., 105
Osaka, T., see Kasai, M., 319

Palacios, J.M., see Cymerman, U., 13
Palmeri, A., see Prato, A., 303
Pan-Hou, H., Suda, Y., Sumi, M., Yoshioka, M. and Kawai, N., Inhibitory effect of 2,4-dihydroxyphenylacetylasparagine, a common moiety of spider toxin, on glutamate binding to rat brain synaptic membranes, 199
Pethő, G., see Barthó, L., 146
Pietra, P., see Angioy, A.M., 263
Polak, J.M., see Inyama, C.O., 111
Prato, A., Scapagnini, U., Rapisarda, E., Giuffrida, R., Palmeri, A., Caruso, A. and Clementi, G., Analgesic activity of a synthetic analogue of eel-calcitonin and its effect on the spontaneous activity of brainstem neurons in the rat, 303

Rapisarda, E., see Prato, A., 303
Reid, J. and McCulloch, J., Capsaicin and blood–brain barrier permeability, 165
Reiter, R.J., Britt, J.H. and Armstrong, J.D., Absence of a nocturnal rise in either norepinephrine, *N*-acetyltransferase, hydroxyindole-*O*-methyltransferase or melatonin in the pineal gland of the domestic pig kept under natural environment photoperiods, 171
Roberts, M.H., see Cassone, V.M., 29
Roisin, M.P., see Anikstejn, L., 291
Rommelspacher, H., see Susilo, R., 325

Saffran, B.N. and Crutcher, K.A., Putative gliotoxin, α-aminoadipic acid, fails to kill hippocampal astrocytes in vivo, 215
Saitoh, T., see Mizuno, Y., 204
Sánchez-Blázquez, P., see Lérida, M., 155
Sandri, C., see Yasargil, G.M., 63
Satoh, M., see Ueda, H., 309
Scapagnini, U., see Prato, A., 303
Schenk, S., Lacelle, G., Gorman, K. and Amit, Z., Cocaine self-administration in rats influenced by environmental conditions: implications for the etiology of drug abuse, 227

Schober, M., see Volknandt, W., 241
Schofield, G.G., see Ikeda, S.R., 123
Sharif, N.A. and Burt, D.R., Guanine nucleotide regulation of receptor binding of thyrotropin-releasing hormone (TRH) in rat brain regions, retina and pituitary, 339
Shen, R.-s., see Awasthi, Y.C., 159
Shioda, S., Nakai, Y., Hori, T. and Sunayama, H., Innervation of thyrotropin-releasing hormone-containing neurons in the bullfrog pars intermedia, 53
Simons, D.J. and Land, P.W., A reliable technique for marking the location of extracellular recording sites using glass micropipettes, 100
Singh, S.V., see Awasthi, Y.C., 159
Skangiel-Kramska, J., see Cymerman. U., 13
Södersten, P., see Forsberg, G., 151
Sone, N., see Mizuno, Y., 204
Sora, I., see Mizukawa, K., 105
Sora, Y.H., see Mizukawa, K., 105
Spyer, K.M., see Numao, Y., 279
Suda, Y., see Pan-Hou, H., 199
Sumi, M., see Pan-Hou, H., 199
Sunayama, H., see Shioda, S., 53
Susilo, R., Damm, H., Rommelspacher, H. and Höfle, G., Biotransformation of 1-methyl-1,2,3,4-tetrahydro-β-carboline-1-carboxylic acid to harmalan, tetrahydroharman and harman in rats, 325
Suzuki, K., see Mizuno, Y., 204
Szabo, T., Blähser, S., Denizot, J.P. and Véron-Ravaille, M., High substance P-like immunoreactivity (SPLI) in the olfactory and electrosensory cells of gymnotid fish, 245
Szolcsányi, J., see Barthó, L., 146

Takagi, H., see Ueda, H., 309
Tokutomi, N., Ikemoto, Y., Nagasawa, K. and Akaike, N., Inhibition of Na^+-dependent GABA uptake in isolated frog sensory nerve cell bodies by extra- and intracellular Li^+, 314
Tomassini Barbarossa, I., see Angioy, A.M., 263
Törk, I., see Mulligan, K.A., 7
Tsuji, S., Minota, S. and Kuba, K., Regulation of two ion channels by a common muscarinic receptor-transduction system in a vertebrate neuron, 139

Ueda, H., Fukushima, N., Ge, M., Takagi, H. and Satoh, M., Presynaptic opioid κ-receptor and regulation of the release of Met-enkephalin in the rat brainstem, 309

Vallat, J.M., see Hugon, J., 1
Véron-Ravaille, M., see Szabo, T., 245
Villalobos, J. and Ferssiwi, A., The differential ascending projections from the anterior, central and posterior regions of the lateral hypothalamic area: an autoradiographic study, 89
Villalobos, J. and Ferssiwi, A., The differential descending projections from the anterior, central and posterior regions of the lateral hypothalamic area: an autoradiographic study, 95
Vodyanoy, V., Muller, D., Kramer, K., Lynch, G. and Baudry, M., Functional reconstitution of *N*-methyl-D-aspartate receptors in artificial lipid bilayers, 133
Volknandt, W., Schober, M., Fischer-Colbrie, R., Zimmermann, H. and Winkler, H., Cholinergic nerve terminals in the rat diaphragm are chromogranin A immunoreactive, 241

Wahnschaffe, U., see Fritzsch, B., 48
Walberg, F., Nordby, T. and Dietrichs, E., The olivonodular projection: a re-examination based on folial cerebellar implants, 82
Weight, F.F., see Ikeda, S.R., 123
Westerman, R.A., Widdop, R.E., Hogan, C. and Zimmet, P., Non-invasive tests of neurovascular function: reduced responses in diabetes mellitus, 177
Wharton, J., see Inyama, C.O., 111
Widdop, R.E., see Westerman, R.A., 177
Wiesenfeld-Hallin, Z., see Forsberg, G., 151
Williams, M., see Jarvis, M.F., 69
Winkler, H., see Volknandt, W., 241
Wu, J.-Y., see Docherty, M., 232

Yamashita, H., see Kasai, M., 319
Yasargil, G.M. and Sandri, C., Morphology of the Mauthner axon inhibitory system in tench (*Tinca tinca* L.) spinal cord, 63
Yoshioka, M., see Pan-Hou, H., 199
Young, A.M.J., see Bradford, H.F., 296

Zahm, D.S., Neurotensin-immunoreactive neurons in the ventral striatum of the adult rat: ventromedial caudate-putamen, nucleus accumbens and olfactory tubercle, 41
Zahniser, N.R., see Miller, J.A., 345
Zeise, M.L., see Knöpfel, T., 188
Zieglgänsberger, W., see Knöpfel, T., 188
Zimmermann, H., see Volknandt, W., 241
Zimmet, P., see Westerman, R.A., 177
Zinebi, F., see Fagni, L., 285